QJM: AN INTERNATIONAL JOURNAL OF MEDICINE

Oxford University Press

<u>QJM</u>. 2020 Mar 30 : hcaa110. Published online 2020 Mar 30. doi: <u>10.1093/qjmed/hcaa110</u> PMCID: PMC7184387 PMID: <u>32227218</u>

Mental health in the Covid-19 pandemic

W Cullen, h1 G Gulati, h2 and B D Kellyh3

^{h1}From the School of Medicine, University College Dublin, Belfield, Dublin 4, Ireland
^{h2} Graduate Entry Medical School, University of Limerick, Limerick, Ireland
^{h3} Department of Psychiatry, Trinity College Dublin, Trinity Centre for Health Sciences, Tallaght University
Hospital, Tallaght, Dublin 24 D24 NR0A, Ireland
Address correspondence to Prof. Walter Cullen, School of Medicine, Health Sciences Centre, University
College Dublin, Belfield, Dublin 4, Ireland. email: walter.cullen@ucd.ie

<u>Copyright</u> © The Author(s) 2020. Published by Oxford University Press on behalf of the Association of Physicians. All rights reserved. For permissions, please email: journals.permissions@oup.com

This article is published and distributed under the terms of the Oxford University Press, Standard Journals Publication Model

(https://academic.oup.com/journals/pages/open_access/funder_policies/chorus/standard_publication_model)

This article is made available via the PMC Open Access Subset for unrestricted re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for the duration of the COVID-19 pandemic or until permissions are revoked in writing. Upon expiration of these permissions, PMC is granted a perpetual license to make this article available via PMC and Europe PMC, consistent with existing copyright protections.

During any outbreak of an infectious disease, the population's psychological reactions play a critical role in shaping both spread of the disease and the occurrence of emotional distress and social disorder during and after the outbreak. Despite this fact, sufficient resources are typically *not* provided to manage or attenuate pandemics' effects on mental health and wellbeing.1 While this might be understandable in the acute phase of an outbreak, when health systems prioritize testing, reducing transmission and critical patient care, psychological and psychiatric needs should not be overlooked during any phase of pandemic management.

There are many reasons for this. It is known that psychological factors play an important role in adherence to public health measures (such as vaccination) and in how people cope with the threat of infection and consequent losses.1 These are clearly crucial issues to consider in the management of any infectious disease, including Covid-19. Psychological reactions to pandemics include maladaptive behaviours, emotional distress and defensive responses.1 People who are prone to psychological problems are especially vulnerable.

All of these features are in clear evidence during the current Covid-19 pandemic. One study of 1210 respondents from 194 cities in China in January and February 2020 found that 54% of respondents rated the psychological impact of the Covid-19 outbreak as moderate or severe; 29% reported moderate to severe anxiety symptoms; and 17% reported moderate to severe depressive symptoms.2 Notwithstanding possible response bias, these are very high proportions—and it is likely that some

Mental health in the Covid-19 pandemic

people are at even greater risk. During the 2009 H1N1 influenza outbreak ('swine flu'), a study of mental health patients found that children and patients with neurotic and somatoform disorders were significantly over-represented among those expressing moderate or severe concerns.<u>3</u>

Against this background, and as the Covid-19 pandemic continues to spread around the world, we hypothesize a number of psychological impacts that merit consideration now rather than later.

In the first instance, it should be recognized that, even in the normal course of events, people with established mental illness have a lower life expectancy and poorer physical health outcomes than the general population. As a result, people with pre-existing mental health and substance use disorders will be at increased risk of infection with Covid-19, increased risk of having problems accessing testing and treatment and increased risk of negative physical *and* psychological effects stemming from the pandemic.

Second, we anticipate a considerable increase in anxiety and depressive symptoms among people who do not have pre-existing mental health conditions, with some experiencing post-traumatic stress disorder in due course. There is already evidence that this possibility has been under-recognized in China during the current pandemic.5

Third, it can be anticipated that health and social care professionals will be at particular risk of psychological symptoms, especially if they work in public health, primary care, emergency services, emergency departments and intensive or critical care. The World Health Organization has formally recognized this risk to healthcare workers,<u>6</u> so more needs to be done to manage anxiety and stress in this group and, in the longer term, help prevent burnout, depression and post-traumatic stress disorder.

There are several steps that can and should be taken now to minimize the psychological and psychiatric effects of the Covid-19 pandemic.

First, while it might be ostensibly attractive to re-deploy mental health professionals to work in other areas of healthcare, this should be avoided. Such a move would almost certainly worsen outcomes overall and place people with mental illness at disproportionate risk of deteriorations in physical *and* mental health. If anything, this group needs enhanced care at this time.

Second, we recommend the provision of targeted psychological interventions for communities affected by Covid-19, particular supports for people at high risk of psychological morbidity, enhanced awareness and diagnosis of mental disorders (especially in primary care and emergency departments) and improved access to psychological interventions (especially those delivered online and through smartphone technologies). These measures can help diminish or prevent future psychiatric morbidity.

Finally, there is a need for particular focus on frontline workers including, but not limited to, healthcare staff. In the USA, the Centers for Disease Control and Prevention offer valuable advice for healthcare workers in order to reduce secondary traumatic stress reactions, including increased awareness of symptoms, taking breaks from work, engaging in self-care, taking breaks from media coverage and asking for help.<u>8</u> This kind of advice needs to be underpinned by awareness of this risk among employers, enhanced peer-support and practical assistance for healthcare workers who find themselves exhausted, stressed and feeling excessive personal responsibility for clinical outcomes during what appears to be the largest pandemic of our times.

Even in this emergency circumstance, or *especially* in this emergency circumstance, we neglect mental health at our peril and to our long-term detriment.

Conflict of interest: None declared.

References

1. Taylor S. The Psychology of Pandemics: Preparing for the Next Global Outbreak of Infectious Disease. Newcastle upon Tyne, Cambridge Scholars Publishing, 2019. [Google Scholar]

2. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al.Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. Int J Environ Res Public Health 2020; 17:1729. [PMC free article] [PubMed] [Google Scholar]

3. Page LA, Seetharaman S, Suhail I, Wessely S, Pereira J, Rubin GJ.. Using electronic patient records to assess the impact of swine flu (influenza H1N1) on mental health patients. J Ment Health 2011; 20:60–9. [PubMed] [Google Scholar]

4. Rodgers M, Dalton J, Harden M, Street A, Parker G, Eastwood A.. Integrated care to address the physical health needs of people with severe mental illness: a mapping review of the recent evidence on barriers, facilitators and evaluations. Int J Integr Care 2018; 18:9. [PMC free article] [PubMed] [Google Scholar]

5. Duan L, Zhu G.. Psychological interventions for people affected by the COVID-19 epidemic. Lancet Psychiatry 2020; 7:300–2. [PMC free article] [PubMed] [Google Scholar]

6. World Health Organization. Mental Health Considerations During COVID-19 Outbreak. Geneva, World Health Organization, 2020.

7. Ho CS, Chee CY.. Ho RC Mental health strategies to combat the psychological impact of COVID-19: beyond paranoia and panic. Ann Acad Med Singapore 2020; 49:1–3. [PubMed] [Google Scholar]

8. Centers for Disease Control and Prevention. *Manage Anxiety & Stress* Atlanta, GA, Centers for Disease Control and Prevention, 2020.